

HYPERAMMONEMIA

When to Suspect a Late-Onset Urea Cycle Disorder (UCD)

Patients with late-onset UCDs may function relatively normally for decades until stressors such as catabolic events, protein overload, or certain drugs trigger a hyperammonemic crisis.^{1,2} Nonspecific symptoms and non-related illnesses experienced by the patient can make it difficult to recognize a late-onset UCD.^{1,2}

A detailed patient/family history and physical examination are essential to identifying patients with a suspected late-onset UCD.^{1,3} If presentation suggests hyperammonemia, check ammonia levels and consider consultation with a metabolic geneticist without delay.¹

Key investigative questions to consider:

- **Is there a family history of metabolic disease, neurological or psychiatric disorders, infant mortality, or consanguinity?**^{1,2}

- **Does the patient's standard diet, recent food intake, or weight changes increase suspicion?**

Symptoms:

- Poorly documented food allergies or intolerances, atypical colic, recurrent vomiting, and/or difficult-to-control gastroesophageal reflux³
- Autoselective vegetarianism (i.e., elective decreased protein intake) or a high carbohydrate intake¹

Triggers:

- Consumption of a high-protein meal (e.g., a barbecue), diet, or supplements^{2,4,5}
- Rapid weight loss (e.g., caused by malabsorption and nutritional disruption following bariatric surgery or extreme mental stress)^{1,5}

- **Does the patient exhibit uncharacteristic behavior or have a history of behavioral or psychiatric illness?**¹

Example: Confusion or combative or psychotic behavior post-partum^{6,7}

- **Does the patient have a history of prolonged clinical courses with seemingly routine illnesses?**¹

Example: Past episodes of flu-like illnesses that resulted in prolonged illness¹

- **Are the patient's nonspecific symptoms unexplained by other differentials?**

Example: Intoxication suspected,² but toxicology screen results are normal

- **Is the patient non-responsive to specific treatment?**

Example: Viral illness suspected to be exacerbating asthma, but response to IV steroids and asthma-specific drugs is negative¹

References

- Summar ML, Barr F, Dawling S, Smith W, Lee B, Singh RH, Rhead WJ, Sniderman King L, Christman BW. Unmasked adult-onset urea cycle disorders in the critical care setting. *Crit Care Clin* 2005;21:S1-S8.
- Haberle J, Boddaert N, Burlina A, Chakrapani A, Dixon M, Huemer M, Karall D, Martinelli D, Sanjurjo Crespo P, Santer R, Servais A, Valayannopoulos V, Lindner M, Rubio V, Dionisi-Vici C. Suggested guidelines for the diagnosis and management of urea cycle disorders. *Orphanet J Rare Dis* 2012;7:32.
- Smith W, Kishnani PS, Lee B, Singh RH, Rhead WJ, Sniderman King L, Smith M, Summar M. Urea cycle disorders: Clinical presentation outside the newborn period. *Crit Care Clin* 2005;21:S9-S17.
- Panlaqui OM, Tran K, Johns A, McGill J, White H. Acute hyperammonemic encephalopathy in adult onset ornithine transcarbamylase deficiency. *Intensive Care Med* 2008;34:1922-1924.
- Collen JF, Nealanjan PD, Koff JM, Neff RT, Abbott KC. Hemodialysis for hyperammonemia associated with ornithine transcarbamylase deficiency. *Appl Clin Genet* 2008;1:1-5.
- Blair NF, Cremer PD, Tchan MC. Urea cycle disorders: a life-threatening yet treatable cause of metabolic encephalopathy in adults. *Pract Neurol* 2015;15:45-48.
- Fassier T, Guffon N, Acquaviva C, D'Amato T, Durand DV, Domenech P. Misdiagnosed postpartum psychosis revealing a late-onset urea cycle disorder. *Am J Psychiatry* 2011;168(6):576-580.